

CORRELATION OF MAP UNITS

Map Unit	Geologic Period	Geologic Epoch	
Qal, Qae, Qp, Qi, Qt	QUATERNARY	Holocene	
Kcb, Kmu, Kmf, Kml, Kd, Klc, Kjc, Kmb, Kk, Kf, Kc, Ks, Kt, Kx			TERTIARY
Kmu, Kmf, Kml, Kd	Paleocene		
Kmu, Kmf, Kml, Kd		Upper Cretaceous	
Kmu, Kmf, Kml, Kd	Lower Cretaceous		
Jmb, Jms, Js, Jt, Jx		JURASSIC	
Jmb, Jms, Js, Jt, Jx	Middle Jurassic		
Tsk, Tsw, Tsc, Txb	TRIASSIC		
Tsk, Tsw, Tsc, Txb		Lower Triassic	
Xb		EARLY PROTEROZOIC	

DESCRIPTION OF MAP UNITS

Qal Alluvium (Holocene)—Unconsolidated clay, silt, sand, and minor gravel deposits on floors of many washes and some canyons. Includes stream-channel and flood-plain deposits, and low-level alluvial-terrace deposits as much as 10 m (33 ft) above present level of stream-channel floors.

Qae Alluvial and eolian deposits (Holocene)—Mostly silt and sand occurring on ridges and mesa tops in southeastern part of map area.

Qp Pediment deposits (Holocene and Pleistocene)—Unconsolidated and semi-consolidated silt, sand, and gravel veneer on pediment surfaces. Semiconsolidated conglomerate and conglomeratic sandstone occurs at base of many pediment deposits.

Qi Landslide deposits (Holocene and Pleistocene)—Chaotically mixed masses of sandstone blocks and silt shale mostly derived from Farrer Formation. Shown only in southwestern part of map area where conspicuously large deposits occur.

Qt Terrace gravel deposits (Pleistocene)—Mostly boulder-size gravel deposited on terraces cut by Colorado River.

Tpau Green River Formation (Eocene)
Parachute Creek Member, upper part—Gray and brown, thin-bedded mudstone and siltstone, gray and brown, very fine grained to medium-grained, thin- to thick-bedded, partly cross-laminated and ripple-marked sandstone, and minor tuff and oolitic limestone. Contact with lower part of Parachute Creek Member placed at base of Horse Bench Sandstone bed about 125 m (410 ft) above Mahogany oil-shale bed. Unit incompletely preserved in map area; maximum preserved thickness about 200 m (720 ft).

Tpdl Parachute Creek Member, lower part—Mostly gray and brown siltstone and very fine grained sandstone, few beds of oil shale, and oolitic and algal limestone. Contact with underlying Douglas Creek Member placed at base of Mahogany oil-shale bed. Thickness approximately 140 m (450 ft) sandstone. Mahogany oil-shale bed—Dark bluish-gray weathering, laminated, and commonly obscured by vegetation or soil in map area. Thickness of Mahogany oil-shale bed less than 1 m (3 ft) in map area.

Tgda Douglas Creek Member
Tongue a—Mostly gray to brown, fine to medium-grained sandstone, gray and green siltstone, few shale beds, and oolitic and algal limestone beds, few oil-shale beds in northern part of area. Contact with underlying Wasatch units placed at uppermost basaltic sandstone or red shale in the Wasatch. Thickness ranges from approximately 65 to 115 m (210 to 375 ft).

Tgdc Tongue c—Mostly green and gray siltstone and shale, brown and gray sandstone, brown and gray algal, oolitic, and oolitic limestone, and few thin beds of oil shale and marlstone. Contact with overlying and underlying units of Renegade Tongue of Wasatch Formation placed at beds of red shale or fluviatile sandstone in the Renegade. Thickness ranges from 0 m (0 ft) in T. 16 S., R. 22 E. to about 60 m (200 ft) in T. 16 S., R. 23 E., and about 90 m (300 ft) in T. 16 S., R. 24 E.; pinchout in subsurface trends northward.

Tgdd Tongue d—Mostly gray shale, and oolitic and algal limestone. Contact with overlying and underlying Wasatch units placed at bed of red shale or fluviatile sandstone in the Renegade. Thickness is approximately 1-2 m (3-7 ft) thick in southern part, as much as 15 m (50 ft) thick in western part, and as much as 30 m (100 ft) thick in eastern part of map area.

Tw Wasatch Formation (Eocene and Paleocene)
Renegade Tongue—Brown and gray, fine to medium-grained, thick-bedded, partly cross-bedded sandstone and shale, and red and gray siltstone and shale. Thickness of Renegade Tongue commonly about 250 m (825 ft), as much as 300 m (1,000 ft) at type locality in T. 17 S., R. 21 E. Unit is and occurs east of center of T. 17 S., R. 22 E. where Douglas Creek Member and Renegade intertongue.

Twu Unit of Renegade Tongue—Mostly medium to thick sandstone, indistinctly bedded, and sparse shale; includes unmapped b tongue of Douglas Creek Member of Green River Formation.

Twv Unit of Renegade Tongue—Mostly red and gray shale; contains large amount of sandstone where it joins main body of Renegade Tongue.

W Wasatch Formation, main body—Dark brown conglomerate and conglomeratic sandstone, containing pebbles of black chert and varcolored quartzite, commonly occurring at base of formation, very light brown and gray, fine to medium-grained, regularly bedded sandstone, and red and greenish-gray silt shale and siltstone, variegated in places. Contact with underlying Tongue of Renegade Tongue of Wasatch Formation placed at base of conglomeratic sequence or lowest red or greenish-gray shale. Thickness ranges from about 1,200 m (3,900 ft) in western part of map area to as little as 125 m (400 ft) in eastern part of map area.

Kt Tucher Formation (Upper Cretaceous)—Mostly brown and gray, fine to medium-grained, commonly thick-bedded sandstone, cross-bedded in most places, and olive to greenish-gray, silt shale. Uppermost sandstone locally laminated, and locally conglomeratic. Contact with underlying Farrer Formation placed at base of succession of thick sandstone units but indistinct in many places. Thickness ranges from approximately 100 m (325 ft) to 200 m (650 ft). Mapped to 109° W. meridian, about 4 km (2.5 mi) east of Utah-Colorado boundary, equivalent unit in Colorado is Hunter Canyon Formation.


Kf Farrer Formation (Upper Cretaceous)—Mostly gray to brown, medium-grained, thin- to thick-bedded, commonly cross-bedded sandstone, greenish-gray, silt shale, and locally, sparse carbonaceous shale beds in lower part. Contact with underlying Neelen Formation gradational, placed where dominantly greenish-gray beds in Farrer grade downward to dominantly carbonaceous beds in Neelen. Thickness ranges from about 125 m (400 ft) to about 250 m (825 ft) in map area. Mapped to 109° W. meridian, about 4 km (2.5 mi) east of Utah-Colorado boundary, equivalent rocks in Colorado included in upper part of Mount Garfield Formation.

Kn Neelen Formation (Upper Cretaceous)—Light brown to brown and light-gray, very fine to fine-grained, flat and cross-laminated to medium-bedded sandstone, medium- to very dark gray carbonaceous shale and silt shale, and small amounts of greenish-gray shale. Sandstone and shale in about equal proportions. Unit contains four coal seams and cross-bedded in western part in extreme southwestern part of map area. Contact with underlying Segó Sandstone usually distinct. Thickness ranges from about 60 m (200 ft) to about 150 m (500 ft). Mapped to 109° W. meridian, about 4 km (2.5 mi) east of Utah-Colorado boundary, equivalent rocks in Colorado included in lower part of Mount Garfield Formation.

Ks Segó Sandstone (Upper Cretaceous)—Very light gray and light-gray to light-brown, fine-grained, flat and cross-laminated to medium-bedded, partly brown, fine-grained, flat and cross-laminated to medium-bedded, and locally conglomeratic sandstone, and sparse medium-gray sandy and silt shale. Shale becomes proportionately more abundant west of R. 20 E. Unit contains unmapped Anchor Mine Tongue of Mancos Shale in eastern part of map area. Contact with underlying Buck Tongue of Mancos Shale gradational, placed where they silt sandstone grades downward to silt shale. Thickness ranges from approximately 45 m (150 ft) to about 60 m (210 ft).

Kmb Buck Tongue of Mancos Shale (Upper Cretaceous)—Medium- to dark-gray shale, silt and sandy in uppermost part, contains sparse limy sandstone lenses, abundant plates of selenite, and carbonized flora. Contact with underlying Castlegate Sandstone abrupt but commonly covered, placed at top of uppermost cuesta-forming sandstone unit. Thickness ranges from approximately 110 m (360 ft) at eastern boundary of map area to approximately 30 m (100 ft) at western boundary.

Kc Castlegate Sandstone (Upper Cretaceous)—Brown to very light gray, very fine to medium-grained, laminated to medium-bedded sandstone, and sparse gray siltstone and shale; contains lenses and pods of sandy and silt marl in eastern part of map area. Unit cross-laminated and cross-bedded in western part of map area, flat laminated and ripple marked in eastern part of map area. Where unit overlies Mancos Shale, gradational and intertonguing contact placed where sandstone grades down to sandy and silt shale; where unit overlies



Base from U.S. Geological Survey, 1988
25000-foot grid rocks based on Utah
contoured from base map, used
Colorado coordinate system, center zone
1000-foot Universal Transverse Mercator
grid, zone 12.

SCALE 1:62,500










1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20				

SCALE 1:94,000

1	2	3	4	5	6	7	8	9	10	11	12
13	14	15	16	17	18	19	20				

INDEX SHOWING TOPOGRAPHIC QUADRANGLES IN MAP AREA

1	2	3	4	5	6	7	8	9	10	11	12
13	14	15	16	17	18	19	20				



GEOLOGIC MAP OF THE WESTWATER 30' x 60' QUADRANGLE, GRAND AND UTAH COUNTIES, UTAH AND GARFIELD AND MESA COUNTIES, COLORADO

By
J. L. Gualtieri
1988

DISCUSSION

Data for the geologic map of the Westwater 30' x 60' quadrangle were derived from published mapping, principally that of Cashion (1967, 1973). Rock unit contacts above the base of the Wasatch Formation were as previously mapped and were cartographically modified to fit the topography of the Westwater base. Likewise, contacts of rock units below the Castlegate Sandstone or Blackhawk Formation were used as previously mapped by Cashion (1973), but were slightly modified in the narrow belt of outcrop including the Dakota Sandstone and Blackhawk Formation. The narrow belt of outcrop including the Dakota Sandstone and Blackhawk Formation was used as previously mapped by Cashion (1973), but was slightly modified in the narrow belt of outcrop including the Dakota Sandstone and Blackhawk Formation. The narrow belt of outcrop including the Dakota Sandstone and Blackhawk Formation was used as previously mapped by Cashion (1973), but was slightly modified in the narrow belt of outcrop including the Dakota Sandstone and Blackhawk Formation.

Mapping in the course of this project was restricted to that part of the section lying between the top of the Mancos Shale and the base of the Wasatch Formation. Reasons of geographic positions of contacts between the several map units are minor but an accurate comparison cannot be made because some mapping, as for example that of Fisher (1936), is on a planimetric base.

The purpose of mapping undertaken in this project was to map the Segó Sandstone and the coal zones in the underlying Neelen Formation, which are shown on other larger-scale maps, the top of the Segó Sandstone is the datum used to correlate the coal zones.

Structure contours were drawn on the top of the Segó and Dakota Sandstone. The Segó was chosen as a structural datum because of its stratigraphic proximity to the overlying coal zones in the Neelen Formation and because its top could be mapped in outcrop and recognized in geophysical logs. Segó structure contours in the subsurface were drawn from data obtained from about 70 petroleum, natural gas, and coal drill holes located in a belt extending from the southwestern part of the map area to the north-central and northeastern parts. No subsurface data were available in the northeastern part; in that area, the location of the contours is inferred from data extrapolated from outcrops along the canyon of the Green River and from geophysical logs of holes drilled north of the northeastern part of the map area.

Structure contours were drawn on top of the Dakota Sandstone because that formation forms conspicuous outcrops and its top can be recognized in geophysical logs. Dakota structure contours in the subsurface were drawn from data obtained from about 125 petroleum and natural gas drill holes located in and just outside the map area. Where the Dakota is eroded, contours are extrapolated by projecting them 350 m (1,148 ft) above the top of the Wingate Sandstone.

Many folds occur in the eastern two-thirds of the map area, roughly delineated by the boundary between R. 19 E. and R. 20 E. They lie on the northward-plunging nose of the Uncompahgne Plateau. The folds are clearly evident in the rocks forming the Book Cliffs and are the cause of the sinuous trace of the cliff-forming outcrops seen on the Westwater 30' x 60' quadrangle map. Extension of these structures and the presence of other structures in the subsurface are based on geophysical data obtained from petroleum and natural gas drill holes in the Cisco Dome and Westwater and San Arroyo anticlines, as well as the Wasatch and Green River Formations. In some folds, structure determined on the top of the Segó Sandstone at depth as related to the structure in surface exposures, is incongruous. This may have resulted from the misinterpretation of the structure, the mislocation of surface rocks, or a combination of both. However, it has been demonstrated by Holmes (1979) that some of the structural incongruity in the Uinta Basin is real; structure contours drawn on the tops of the Green River and Green River Formations diverge because of variations in thickness of the Green River Formation.

Faults associated with the folds in the eastern two-thirds of the map area are high-angle (from vertical to 60°) normal faults that resulted from the relaxation of tectonic stress. The Thompson and Salt Valley anticlines and associated synclinal structures in the southwestern part of the map area lie in the region of the Paradox Basin. The structures originated as early as Late Permian time in response to the flow of salt from the Paradox Member of the Hermosa Formation to tectonically controlled basins. Salt flow and contemporaneous formation of salt structures continued through Middle to Early Jurassic time, when salt flow ceased. In Late Cretaceous time the region was tectonically deformed, accentuating the already-existing salt-cored structures. This is evident in the Book Cliffs, along the trace of the structural axis of the Salt Valley anticline, where deformation is recognizable in beds of the Upper Cretaceous Farrer Formation but not in the basal, lower Paleocene beds of the Wasatch Formation.

All faults associated with the Thompson and Salt Valley anticlines are high-angle normal faults and are the result of subsidence following the exsolution of salt.

REFERENCES CITED

Cashion, W. B., 1967, Geology and fuel resources of the Green River Formation, southern Uinta Basin, Utah and Colorado. U.S. Geological Survey Professional Paper 548, p. 1-125.

—, 1973, Geologic and structure map of the Grand Junction quadrangle, Colorado and Utah. U.S. Geological Survey Miscellaneous Geologic Investigations Map I-736, scale 1:250,000.

Fisher, D. J., 1936, The Book Cliffs fold in Emery and Grand Counties, Utah. U.S. Geological Survey Bulletin 852, p. 9, 10, and 11.

Holmes, W. F., 1979, Maps showing generalized structure contours on top of the Wasatch and Green River Formations, geologic sections, and contours of thicknesses of the Green River Formation, southeastern Uinta Basin, Utah and Colorado. U.S. Geological Survey Miscellaneous Investigations Series Map I-1156, scale 1:125,000.